

Land surface Interactions with the Atmosphere over the Iberian Semi-arid Environment (LIAISE): 1st modelling intercomparison

M.A. Jiménez¹, J. Cuxart¹, A. Grau¹,

(1) Universitat de les Illes Balears, Palma, Spain

A. Boone², S. Donier², P. Le Moigne²,

(2) CNRM, Meteo-France/CNRS, Toulouse, France

J.R. Miró³, Jordi More³,

(3) Servei Meteorològic de Catalunya, Barcelona, Spain

A. Tiesi⁴, P. Malguzzi⁴,



(4) CNR-ISAC, Bologna, Italy

J. Brooke⁵ and Martin Best⁵




(5) Met Office, Exeter, UK





The 1st mesoscale intercomparison case



17th July 2016 at 1200 UTC



El Poal (V8)



The 1st mesoscale intercomparison case

16-18 July 2016

* clear skies, A conditions

* locally/basin/mesoscale generated winds (interact)


Models

MesoNH (MNH)

MOLOCH (MOL)

Unified Model (UM)

WRF



Model setup

- **36h run** (from 16 July at 1800 UTC to 18 July at 0600UTC)
- **2 nested domains** (1-way): 2km x 2 km and 400m x 400m resolution (540 x 450 gridpoints)
- **Vertical resolution** (2m and stretched above, 85 levels)
- **Initial/Lateral BC:** ECMWF
- **Differences:** Turbulence, Radiation (5min), Surface


10 m (agl)
wind vectors
(MesoNH)


17th July
0600 UTC

E-wind prevail


17th July
1500 UTC

SB front
interacts with
local winds







El Poal (V8) - FLOOD



Validation using AWS

Moloch WRF MesoNH UM



Mean BIAS (model-obs)

Validation using AWS

MesoNH

Moloch


UM

WRF


ALL run,

DAY (1000-1400 UTC), NIGHT (0000-0400 UTC)


Wind speed: about +0.5m/s




Temperature: about +1.5°C




Wind direction: [-60°, +30°]








Specific humidity: about -1g/kg



Averaged temperature over the regions








✓ Spatial resolution
 ✓ Soil features






Validation using LST

Surface thermal gradient




Validation using LST



Variability of SEB

SEB terms averaged for each region (drip irrigated)

All the models do not present significant differences in the SEB terms for each region




Ground Flux

Latent Heat Flux

Sensible Heat Flux

Net Radiation


Variability of the surface cover




- ✓ Models present differences in the surface parameters (LAI, fveg, albedo, ...)

Variability of the surface cover


MesoNH




UM



WRF



MOLOCH



Summary

- The case 16-18 July 2016 is taken for the 1st mesoscale intercomparison.
- ✓ **Locally-generated circulations**
(interaction between local, basin, mesoscale)
- **Results (known features):** models are able to reproduce the general patterns of the region **BUT:**
 - ✓ **Models tend to overestimate wind speed** (daytime)
 - ✓ **Difficulties in reproducing nocturnal nearly calm conditions**
 - ✓ **Temperatures are overestimated** (specially during night-time)

Summary

- Models are not able to reproduce the heterogeneities:
 - ✓ **Surface model** (processes included, irrigation)
 - ✓ **Surface parameters & initialization**
(irrigated, rainfed,... zones)
 - ✓ **Parameterizations** (turbulence, advection, radiation)
- **Sensitivity tests (work in progress)**
 - ✓ **Initial and lateral BC** (GFS, NCEP)
 - ✓ **Surface features**
(soil moisture, vegetation, surface model...)
 - ✓ **Spatial resolution**
- **After testing models + LIAISE campaign:** possible future
(GEWEX) intercomparison based on IOPs?