

# On the relevance of diurnal cycle model improvements to weather & climate extremes

Gianpaolo Balsamo

ECMWF, Earth System Modelling Section, Coupled Processes Team

[gianpaolo.balsamo@ecmwf.int](mailto:gianpaolo.balsamo@ecmwf.int)


Aknowledgements: ECMWF, Copernicus, GEWEX

**2018 GEWEX Open Science Conference, Canmore, Alberta, Canada (Planet-A)**





# Coupling for improved diurnal cycle and impact on extremes



- **Introduction: the ECMWF forecasting systems in 2018**
- Three examples for surface-atmosphere coupling relevance:
  - Ocean-coupling effects on diurnal cycle of temperature and cyclones
  - Soil-coupling effects on soil moisture and extreme surface temperature
  - Snow-coupling effects on snow depth and extreme surface temperature




5-10<sup>th</sup> July 2014



2-10<sup>th</sup> Aug 2017



20-28<sup>th</sup> Feb 2018



- Outlook
  - Improved surface coupling as key to hydrological & environmental applications

# Earth surface modelling components @ECMWF in 2018

## • NEMO3.4


NEMO3.4 (Nucleus for European Modelling of the Ocean)

[Madec et al. \(2008\)](#)

[Mogensen et al. \(2012\)](#)

ORCA1\_Z42: 1.0° x 1.0°

ORCA025\_Z75 : 0.25° x 0.25°



## • EC-WAM


ECMWF Wave Model

[Janssen, \(2004\)](#)

[Janssen et al. \(2013\)](#)

ENS-WAM : 0.25° x 0.25°

HRES-WAM: 0.125° x 0.125°



## • LIM2


The Louvain-la-Neuve [Sea Ice Model](#)

[Fichefet and Morales Maqueda \(1997\)](#)

[Bouillon et al. \(2009\)](#)

[Vancoppenolle et al. \(2009\)](#)

ORCA025\_Z75 : 0.25° x 0.25°



## • Hydrology-**TESSEL**

[Balsamo et al. \(2009\)](#)  
[van den Hurk and Viterbo \(2003\)](#)

Global Soil Texture (FAO)

New hydraulic properties

Variable Infiltration capacity & surface runoff revision

## • **NEW SNOW**

[Dutra et al. \(2010\)](#)

Revised snow density

Liquid water reservoir

Revision of Albedo and sub-grid snow cover

## • **NEW LAI**

[Boussetta et al. \(2013\)](#)

New satellite-based

Leaf-Area-Index

## • **SOIL Evaporation**

[Balsamo et al. \(2011\),](#)

[Alberroel et al. \(2012\)](#)

## • **H<sub>2</sub>O / E / CO<sub>2</sub>**

Integration of

Carbon/Energy/Water

[Boussetta et al. 2013](#)

[Agusti-Panareda et al. 2015](#)

## • **Lake & Coastal area**

[Mironov et al \(2010\),](#)

[Dutra et al. \(2010\),](#)

[Balsamo et al. \(2012, 2010\)](#)

Extra tile (9) to for sub-grid lakes and ice

LW tiling (Dutra)





## • **Enhance ML**

Snow ML5

Soil ML9

[Dutra et al. \(2012, 2016\)](#)

[Balsamo et al. \(2016\)](#)



Atmos Land Resol.	ECMWF in 2018
80 km	ERA1
32 km	ERA5+ SEAS5+*
18 km	ENS+*
9 km	HRES+*

## \*Ocean

used across forecast systems and in Ocean reanalysis


(\*migration completed with HRES-coupled operational from the 5th June 2018)

## +Land

used across forecast systems and new Climate reanalysis

# Seamless surface-atmosphere coupling of Integrated Forecasting System



As operational on June 5<sup>th</sup>, 2018



See Buizza et al (2018), Keeley et al (2018), Mogensen et al (2018), published in the ECMWF 2018 Summer Newsletter



# Coupling for improved diurnal cycle and impact on extremes

- Introduction: the ECMWF forecasting systems in 2018
- **Three examples for surface-atmosphere coupling relevance:**
  - Ocean-coupling effects on diurnal cycle of temperature and cyclones
  - Soil-coupling effects on soil moisture and extreme surface temperature
  - Snow-coupling effects on snow depth and extreme surface temperature




- Outlook
  - Improved surface coupling as key to hydrological & environmental applications

# Ocean-coupling and local effects on sea surface temperature (SST)





- The ECMWF Ocean-coupled (red) model is better simulate the cool wake after the passage of Tropical cyclone Neoguri. A more realistic response is observed comparing the 10-day forecast with an on-track DRIBU observation of SST, both for TC passage and diurnal cycle





# Impact of Ocean-coupling along the track of Tropical Cyclone Neoguri

## Sea surface temperature



## Surface heat flux




ECMWF  
Ocean  
Coupled  
5-day  
Forecast

ECMWF  
Ocean  
Uncoupled  
5-day  
Forecast

- Coupled Forecast:
- Gets the SST cooling better
  - Reduced Heat flux to atmos.
  - Gets the TC Intensity Better


# Comparing forecasts with satellite-based sea surface temperature

## Sea surface temperature (forecast)




ECMWF  
Ocean  
Coupled  
5-day  
Forecast

ECMWF  
Ocean  
Uncoupled  
5-day  
Forecast



## Sea surface temperature (observation)



Satellite Observations  
from AMSR2 MW SST


After the passage of  
Tropical Cyclone Neoguri,  
10<sup>th</sup> of July 2014

- Coupled forecast:
- Gets the SST cooling after the passage of Tropical Cyclone in better agreement with EO data of Satellite SSTs



# Impact of Ocean-coupling on Tropical Cyclones and relevance for 2018 season

Tropical Cyclones Intensity is generally improved when looking at recent cases (past 2-years)



The red curve is for the ECMWF HRES Coupled as implemented on the 5<sup>th</sup> of June 2018


Mean-sea-level-pressure, MSLP in hPa, of new 45r1 (red) & 43r3 (blue). The data sample includes about 750 cases at initial time, decreasing to about 200 at forecast day 5-6 and to about 50 at day 10. Bars indicate 95% confidence.

# What happens to the temperature diurnal cycle enhancing surface coupling?

- Towards more realistic surface temperature (skin and below) particularly in clear/sky
- Towards increased variability and surface responsiveness to atmospheric forcing

## *Ocean skin*


Ampl. 1st harmonic diurnal c.; Tskin(C), Month:20160700, Exp:gn6y-0001



Difference in diurnal cycle amplitude due ocean-coup

## *Land skin*



Ampl. 1st harmonic diurnal c.; Tskin(C), Month:20160800, Exp:gqx3-gqx4



Difference due to enhance multi-layer land-coup

# Coupling for improved diurnal cycle and impact on extremes

- Introduction: the ECMWF forecasting systems in 2018
- **Three examples for surface-atmosphere coupling relevance:**
  - Ocean-coupling effects on diurnal cycle of temperature and cyclones
  - Soil-coupling effects on soil moisture and extreme surface temperature
  - Snow-coupling effects on snow depth and extreme surface temperature



- Outlook
  - Improved surface coupling as key to hydrological & environmental applications

# Increased soil model vertical resolution to improve use of satellite data

An enhanced soil vertical layer is motivated by land data assimilation as it shown to better correlate with satellite products of soil moisture.


4-layers:

# 0-7 cm


# 7-28 cm

# 28-100 cm

# 100-289 cm



7 cm



9-layers:

# 0-1 cm

# 1-3 cm

# 3-7 cm

# 7-15 cm



# 15-25 cm

# 25-50 cm

# 50-100 cm

# 100-200 cm

# 200-300 cm



Comparison with ESA-CCI soil moisture remote sensing (multi-sensor) product.(1988-2014). A finer soil model improves the correlation with measured satellite soil moisture

Globally Improved match to satellite soil moisture (shown is Anomaly correlation  $\Delta$ ACC calculate on 1-month running mean)



Thanks to Clément Albergel, Patricia De Rosnay, LDAS-Team


See Dorigo et al. (2017 RSE)

# Impact of the soil model vertical resolution: heatwaves severity

During summer 2017 the effect of multi-layer is examined for European heatwave, here shown for Corboba (Spain) where temperatures went above 40° Celsius on the 6<sup>th</sup> of August 2017



ECMWF  
Land  
model  
**ML9** &  
**ML4**  
(offline)

Difference  
ML9-ML4  
soil model  
(offline)



Extreme heat warnings across southern Europe as temperatures hit 40C and above

Not dangerous Potentially dangerous  
Dangerous Very dangerous





Differences in the maximum skin temperature ML9-ML4

An enhanced soil vertical discretisation is increasing the amplitude of the diurnal cycle. Extremes heatwave are up to 1-2 K hotter

# Coupling for improved diurnal cycle and impact on extremes


- Introduction: the ECMWF forecasting systems in 2018
- **Three examples for surface-atmosphere coupling relevance:**
  - Ocean-coupling effects on diurnal cycle of temperature and cyclones
  - Soil-coupling effects on soil moisture and extreme surface temperature
  - Snow-coupling effects on snow depth and extreme surface temperature



- Outlook
  - Improved surface coupling as key to hydrological & environmental applications


# Increased snow model vertical resolution: impact in cold regions climate

Increased vertical discretization of the snowpack (**up to 5 layers**) permits a better physical processes representation




Difference ML- SL in Snow depth RMSE winter (DJF)

Difference ML - SL in  $T_{skin}$  minimum winter (DJF)



An improved snow depth (ML – SL) evaluated with in-situ SYNOP snow depth. RMSE of 0.19m (0.23m) in ML (SL). This is 17% RMSE error reduction in snow depth.




Winter reduction of the 2m minima temperatures with increasing diurnal-cycle. DIFF Tmin 2-4 K colder in ML compared to SL snow. Increased variability




# Impact of snow model vertical resolution increase on near surface temperature

Increased vertical discretization of the snowpack (**up to 5 layers**) permits a better 2-m forecast: here hourly day-2 forecasts are shown for 24-hour to 47-hour ahead, concatenated to form a continuous time-series



## T2m Observations, T2m forecast (current snow, SL), T2m forecast (ML)



In clear-sky the MULTI-layer snow scheme is capable to produce stronger winter inversions improving observation match.

**NOLIM** indicates a stability limiter safety is deactivated.


The increased variability in the diurnal cycle is beneficial for ensemble forecasting.



# Earth Observations for surface coupled model development: the example of LSTs

Land Surface Temperatures (LSTs) provides validation and guides diurnal cycle model development:

- LST to evaluate impact of vegetation modelling (see Trigo et al. 2015, used EUM-LSA-SAF )
- LST to constrain HTESSEL coupling parameters (see Orth et al., 2017, used ESA-CCI)
- LST to show value multi-layer snow over Antarctica (see Dutra et al 2017, used MODIS)



# Summary

- Model development on diurnal cycle improvements connects to capability of better representing Extreme events (good argument in favor of seamless approach)
- Three cases shown: Ocean-coupling, Soil-layers, Snow-layers enhancements
- Systematic model errors when reduced will introduce larger variability (skill<sub>vs</sub> rmse)
- Energy & Water cycle improvements (e.g. soil moisture, snow) support hydro-apps
- Carbon cycle forecasting skills strongly depend on temperature & moisture, fluxes

## Outlook:

- GEWEX actions linking across-scales (e.g. WWRP NWP/S2S with WCRP/NTCP/X)
- EO-observations (Tskin, MW) can validate increase complexity in surface modelling
- Progressive inclusion across-models of human impact (CO<sub>2</sub>, Water-use, Land-use)

An example in the CO<sub>2</sub> Human Emissions project <https://www.che-project.eu>



# Thank you for your attention

*This work would NOT be feasible & fun without amazing individuals & teams contributions.  
A big thanks to All collaborators!*

**2018 GEWEX Open Science Conference, Canmore, Alberta, Canada (Planet-A)**



# Earth surface monitoring & forecasting advance needs better use EO observations



## remote sensing

Special Issue:

"Advancing Earth Surface Representation via Enhanced Use of Earth Observations in Monitoring and Forecasting Applications"

[http://www.mdpi.com/journal/remotesensing/special\\_issues/earthsurface\\_RS](http://www.mdpi.com/journal/remotesensing/special_issues/earthsurface_RS)

There is time up to 31 July 2018. Discussions at: 2<sup>nd</sup> International Surface Working Group (ISWG) To be hosted by IPMA, Lisbon, 26-28 June 2018.


Hydrol. Earth Syst. Sci., 21, 2483–2495, 2017  
 www.hydrol-earth-syst-sci.net/21/2483/2017/  
 doi:10.5194/hess-21-2483-2017  
 © Author(s) 2017. CC Attribution 3.0 License.



## Advancing land surface model development with satellite-based Earth observations


Rene Orth<sup>1,2</sup>, Emanuel Dutra<sup>1,3</sup>, Isabel F. Trigo<sup>3,4</sup>, and Gianpaolo Balsamo<sup>1</sup>

Table 2. Summary of perturbed model parameters and their characteristics (adapted from O16).



# Extended-range predictability – Connecting the time-scales

- The WCRP Grand Challenge on Near-Term Climate Prediction (NTCP) is designed to link the scales from Annual to Decadal (A2D). This GC is led by A. Scaife and Y Kushnir (Co-Leads)



19 international members of the group

Outcome report (yearly) for the:  
**Global Annual to Decadal  
Climate Update**




# Modelling surface heterogeneity and coupling with the atmosphere

- The processes that are most relevant for near-surface weather prediction are also those that are most interactive and exhibit positive feedbacks or have key role in energy partitioning



**Over Land/Biosphere**

- Snow-cover, ice freezing/melting have positive feedback via the albedo
- Vegetation growth and variability interact with turbulence & moisture
- Vertical heat transport in soil/snow



**Over Ocean/Cryosphere**

- Transition from open-sea to ice-covered conditions
- Sea-state dependent interaction wind induced mixing/waves
- Vertical transport of heat



**Over Water-bodies**

- Lakes have large thermal inertia
- Different albedo & roughness

Spatial heterogeneity calls for high-resolution horizontal/vertical to represent the surface-atmosphere coupling